



HUNTER RIVER CO.

MOXISTAR

LONG ACTING INJECTION FOR CATTLE

PREMIUM CHOICE FOR TICK MANAGEMENT

PERSISTENT CONTROL OF CATTLE WORMS

NO KNOWN EFFECT ON DUNG BEETLES



500ML

500 mL

ACTIVE CONSTITUENT:
100 MG/ML MOXIDECTIN

Moxistar Long Acting Injection has been specially formulated to provide a prolonged release of moxidectin in cattle. It is ideal for long term protection against key internal and external parasites including cattle tick. With persistent activity against internal parasites for up to 120 days (see below table for details) or cattle tick for at least 51 days, no other product offers longer term protection ensuring optimum health, performance and fertility.

MOXISTAR LONG ACTING INJECTION FOR CATTLE is effective against sensitive strains of the following parasites:

Internal parasites: Mature (adult) and immature (L4): Barber's pole worm, *Ostertagia ostertagi* / *Ostertagia lyrata* (including inhibited larvae), Black scour worms, Small intestinal worm, Nodule worm, Hookworm, Whipworm, Lungworm and Adult nematodes.

External parasites: Lice (Sucking lice). Aids in the control of *Bovicola bovis* (biting lice), Mites and Ticks.

PROTECTION PERIOD

When MOXISTAR LONG ACTING INJECTION FOR CATTLE is used at the recommended dose rate as a single subcutaneous injection, it prevents re-infection of cattle with parasites as in the following table:

Parasite Species	Persistent Protection Period
<i>Ostertagia</i> spp.	112 days
<i>Haemonchus</i> spp.	120 days
<i>Trichostrongylus axei</i> .	72 days
<i>Cooperia</i> spp.	21 days
<i>Dictyocaulus viviparus</i>	120 days
<i>Oesophagostomum radiatum</i>	120 days
<i>Linognathus vituli</i>	133 days

APPLICATOR

5ml Vaccinator

WITHHOLDING PERIODS

MEAT: 56 days.

MILK: DO NOT USE in lactating cows or within 80 days of calving where milk may be used or processed for human consumption.

ESI: 108 days. DO NOT USE in cattle that may be exported live.

STORAGE

Store below 30°C (room temperature). Protect from light.


COMPARABLE PRODUCT

Virbac Cydectin Long Acting Injection for Cattle

DOSING & ADMINISTRATION

Dose Rate: 1 mg moxidectin/kg liveweight (1 mL/100 kg liveweight)

INJECT ONLY by subcutaneous injection into the back of the ear (as directed below). Do not inject anywhere else in the animal.



Weight Range (kg)	Total Dose Volume (mL)	No. of Cattle Treated	Weight Range (kg)	Total Dose Volume (mL)	Dose Volume Per Ear (mL)	No. of Cattle Treated
* 100	1.0	500	501-550	5.5	2.75	90
101-150	1.5	333	551-600	6	3	83
151-200	2.0	250	601-650	6.5	3.25	76
201-250	2.5	200	651-700	7	3.5	71
251-300	3.0	166	701-750	7.5	3.75	66
301-350	3.5	142	751-800	8	4	62
351-400	4.0	125	801-850	8.5	4.25	58
401-450	4.5	111	851-900	9	4.5	55
451-500	5.0	100	901-950	9.5	4.75	52
			951-1000†	10	5	50

* DO NOT USE IN CATTLE LESS THAN 100 KG

† DO NOT USE IN CATTLE GREATER THAN 1000 KG

For more information please visit hunterriverco.com.au

Virbac & Cydectin are registered trade marks of Virbac (Australia) Pty Ltd.

