

blank page

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent 

Pilferer 500

HERBICIDE

ACTIVE CONSTITUENT: 500g/L PROPYZAMIDE

GROUP 3 HERBICIDE

For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table.

IMPORTANT: Read this booklet before use.

APVMA Approval No: 68267/122248

AIRR APPARENT PTY LTD
15/16 Princes Street, Newport NSW 2106
P 03 5820 8400 | E enquiries@apparentag.com.au | www.apparentag.com.au

219

DIRECTIONS FOR USE

Restraints:

Do not apply more than once per season.

Do not apply to water logged soils.

Do not apply to land where runoff to aquatic areas or sensitive crops like cereals is likely.

Crops	Weeds Controlled	State	Rate L/ha
Canola	Annual Ryegrass, Winter Grass, Barley	All States	1.0
Grain Legumes (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)	Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue		
Legume Pastures (including lucerne, clover, medics etc. grown for forage, hay or seed production)		NSW, ACT, Tas, SA, WA only	1.0-1.5
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, Seedling Sorrel, Blackberry Nightshade, Patterson's Curse, Seedling Wireweed		1.5-2.0
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All States	4.5 in 330-550L of water
Lettuce (transplanted)		Vic only	2.3 in 300-550L of water
Turf and lawns Common Couch	Winter Grass	All States	1.2 (12mL/100m ²)
Queensland Blue Couch		Qld only	0.6 (6mL/100m ²)
Buffalo Grass		WA only	1.2 (12mL/100m ²)

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

Critical Comments	
Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that move treated soil back into the furrow.	
Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. Do not use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.	
Apply when weeds are at post or pre-emergent stages.	
SEEDLING CROPS: Apply after the third trifoliolate leaf is fully developed.	
ESTABLISHED CROPS: Apply any time before the end of July.	
Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Apply on light sandy soils, low in organic matter.	
Rate/Bowling Green	Apply during the autumn and winter if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld Blue Couch after 4 weeks.
37m x 37m	
170mL	
85mL	DO NOT RESEED FOR 60 DAYS AFTER TREATMENT.
170mL	Use ONLY on established turf.
	DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.

WITHHOLDING PERIODS:

CANOLA:

HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.

GRAIN LEGUMES:

HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.

ALL OTHER CROPS:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of Apparent Pflifer 500 Herbicide to water the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

COMPATIBILITY

Apparent Pflifer 500 Herbicide is compatible with a range of knockdown and pre-emergent herbicides and insecticides. The following table provides some recommendations on the suitability of tank mix combinations.

Physical compatibility with Apparent Pflifer 500 Herbicide should be determined prior to mixing with a product not listed in this table, or when mixing Apparent Pflifer 500 Herbicide as a component of a 3-way tank mix.

Always read the product label for the manufacturer's tank mix recommendations to determine individual product compatibility options and correct mixing orders for individual products. If unsure, perform a jar test before proceeding to determine physical compatibility. Physical compatibility does not always guarantee biological compatibility.

Herbicides	Compatibility recommendation
Apparent Atrazine 900 WG	Compatible
Apparent Diuron 900 WG	Compatible
Ecopar (pyraflufen-ethyl)	Compatible
Glyphosate present as the potassium (K) salt-based products**	Not recommended - incompatible
Apparent Glyphosate 450	Conditional
Glyphosate Dry WG680	Compatible
Apparent Haloxifop 520	Compatible
Apparent Clobbertime 400	Conditional
MCPA products	Not recommended - incompatible
Apparent Oxyflurofen 240 EC	Conditional
Apparent Paraquat 250 SL	Not recommended - incompatible
Paraquat 360 SL	Not recommended - incompatible
Apparent Disturb 875 WG	Compatible
Apparent Trifluralin 480 EC	Not recommended - incompatible

Insecticides	Compatibility recommendation
Dimethoate 400 EC	Not recommended - incompatible
Omethoate 290 SL	Compatible
Apparent Annihilate 100	Compatible
Apparent Stockade 250 EC	Conditional
Trojan (gamma-cyhalothrin)	Compatible

Explanatory notes on recommendations

Compatible: Products considered physically compatible with Apparent Propyzamide 500 WG in a two-way mix.

Conditional: Tank mixes of Apparent Propyzamide 500 WG with the following product may result in nozzle or filter blockage if vigorous agitation in the spray mix unit is not maintained during the entire spray operation.

Not recommended - incompatible: Tank mixes of Apparent Propyzamide 500 WG with the following product is likely to form a slimy residue which can block spray nozzles and filters during application.

** It is not recommended to mix Apparent Propyzamide 500 WG with any products containing ***Glyphosate present as the potassium (K) salt.***

Information on compatibility is understood to be correct at the time of publication, however products may vary from time to time.

WEED RESISTANCE WARNING

GROUP 3 HERBICIDE

Apparent Pflifer 500 Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group 3 herbicide. Some naturally occurring weed biotypes resistant to Apparent Pflifer 500 Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Pflifer 500 Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Pflifer 500 Herbicide to control the resistant weeds.

Advice as to strategies and alternative treatments that can be used, should be obtained from your local supplier,

consultant, local Department of Agriculture, Primary Industries Department or an AIRR Apparent representative.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Safety Directions

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A material safety data sheet for Apparent Pilferer 500 Herbicide is available from AIRR Apparent Pty Ltd on request.

CONDITIONS OF SALE

AIRR APPARENT Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on APPARENT's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of AIRR APPARENT Pty Ltd has any authority to add to or alter these conditions.